

Impact of Oxidative Stress on Male Reproduction: Amelioration by Melatonin and Some Selected Food-Grade Antioxidants

Seema Rai^{1*}, Sunita Patel¹, Vijay K. Bharti² and C. Haldar³

¹Department of Zoology, School of Studies in Life Science, Guru Ghasidas Vishwavidyalaya, Bilaspur – 495009, Chhattisgarh, India; seemarai.72@ggu.ac.in, mesunita8696@gmail.com

²DRDO-Defence Institute of High Altitude Research (DIHAR), Leh – 194101, Ladakh, India; vijaybharti.dihar@gov.in

³Pineal Research Lab, Department of Zoology, Institute of Science, Banaras Hindu University, Varanasi – 221005, Uttar Pradesh, India; chaldarbhu@gmail.com

Abstract

Infertility is a common issue. In India, it has doubled in prevalence, raising health issues in both individuals and communities. Couples who are infertile suffer emotionally, socially, and financially. The Indian Society of Assisted Reproduction estimates that there are up to 27.5 million infertile persons living in India, both men and women. Endocrine disruptors, environmental stress, and abnormalities/imbances in the body's natural antioxidant defence mechanism all put male fertility at greater risk. Studies on how high altitude affects male fertility are very few. Recent research has shown that oxidative stress from different sources impairs spermatogenesis and causes an imbalance in the level of male hormones. There are two primary causes of oxidative stress: extrinsic and intrinsic, responsible for free-radical generation. There is an internal antioxidant defence mechanism that scavenges the reactive oxygen species, i.e., free radical generation which neutralizes oxidative stress. These antioxidants are important for the protection of cellular integrity. Apart from the antioxidants, various food supplements like melatonin, vitamin C, vitamin E, carotenoids, cysteines, etc., are suitable antioxidants for improving male fertility. Therefore, there is a great need for information on some clinically examined edible phyto-oxidants, including melatonin, for amelioration of oxidative stress-induced male infertility. This review focuses on the information available as of now about free radical (ROS)-induced reproductive damages in the plains and the high-altitude regions and the role of various antioxidants, including melatonin, in male infertility.

Keywords: Antioxidant, Male Infertility, Melatonin, Oxidative Stress, Spermatozoa.

1. Introduction

Nearly 30 million men nationwide suffer infertility, ranging between 2.5% and 15% globally^{1,2}. Males account for about 40% of instances of infertility, and several diseases, including varicocele, cryptorchidism, hypogonadism, and hereditary factors, are known to play a role in the infertility process. Furthermore, numerous psychological, physiological, and sociocultural issues have been connected to infertility³. So, it is necessary to understand the mechanism and know the important factors of infertility in males to decipher the key stages and molecules for clinical management. In today's

scenario, the terms oxidative stress, oxidative damage, free radicals load, and antioxidants are the most common terminologies in scientific research and have become an integral part of the scientific debate. Various studies revealed that some of the fundamental and common factors causing male infertility are cellular oxidative damage and free radicals mainly Reactive Oxygen Species (ROS). There is plenty of evidence suggesting that free radicals (oxidative stress) are related to male infertility by reducing sperm motility, sperm DNA damage, decreased sperm production, and various genetic disorders. So, it's critical to understand how cellular oxidative damage is linked with male infertility.

*Author for correspondence

Reproduction is an energy-demanding process⁴. The expense of reproduction stands as a fundamental concept in the realm of evolution and for the perpetuation of the species⁴. No doubt that reproduction requires energy⁵, and to fulfill the energy needs, both basal and local metabolic rates must be enhanced. It is, therefore, suggested that the role of stress in reproduction is inversely proportional to the fertility rate of both males and females. Reproduction contributes to several other activities such as an increase in BMR, and a decrease in immunity and carbohydrate metabolism in several species⁶⁻⁹. Consequently, reproduction enhances energy expenditure by uplifting basal metabolic rate¹⁰. The higher rate of metabolism results in increased free radical production. Therefore, one might expect that reproduction increases vulnerability to oxidative stress; and it is also expected that reproduction will strengthen the body's antioxidant defence mechanisms.

The major target in reproduction affected by oxidative stress and free radicals in a variety of species is the membrane of spermatozoa (lipid peroxidation) thus impairing the motility of sperm¹¹⁻¹³. So, one would expect that the effort of breeding can generate free radicals (oxidative stress) whose efficacy and forte would rely on the effectiveness and accessibility of the redox (antioxidant) defence system¹⁴. As for additional assessments to be carried out for the management and treatment of male infertility, opinions differ concerning which individuals should undergo testing for OS. Furthermore, there is not much information on an antioxidant-based therapeutic regime for male infertility. Meanwhile, the types, doses, and duration of antioxidant action for patients having extreme levels of free radicals in blood and semen are still controversial. Therefore, the present review aims to provide an update on indications about ROS, i.e., free radical production, and induction of oxidative stress associated with male infertility, its immediate/long-term therapeutic control, etc.

2. Role of Oxidative Stress in Male Reproduction

Due to the fast change in lifestyle opted by many people, dependency on processed cuisine, and exposure to a diverse array of chemicals and drugs, etc., play an important role in inducing oxidative stress. In the metabolic system, steady-state redox balance is maintained at a set value, and deviations from it lead to

initiating oxidative stress¹⁵. In other words, when there is an imbalance between the generation of free radicals (ROS) and the body's antioxidant compounds it leads to the induction of oxidative stress^{16,17}. ROS are very erratic and highly reactive molecules, that bind through a diverse group of biomolecules, for example, DNA, proteins, lipids, and a few of the carbohydrates¹⁸⁻²⁰. Hence, Leydig cells, spermatozoa, etc., are vulnerable to ROS thereby affecting male reproductive physiology. The body's antioxidant defence system comprises endogenous and exogenous antioxidants that scavenge ROS and diminish their toxic effect on the cellular system²⁰.

The inflammation in testicular tissues causes infiltration of leukocytes and the production of immature sperm; these are other causes of ROS in human sperm^{21,22}. ROS are produced in the process by which leukocyte activation and chemotaxis promote infection and inflammation of the male reproductive organs. The breakdown of pathogens triggers the leukocyte myeloperoxidase system, which produces ROS²³. The excessive generation of ROS by leukocytes can ultimately result in oxidative stress in semen. These endogenous ROS are the primary factor in the production of abnormal and immature sperm²⁴. Furthermore, cytoplasm deposits in the mid-piece fall off to cause cell elongation and compression²⁴.

There are mainly two different sources of oxidative stress - exogenous and endogenous. Exogenous sources are ionizing radiation, ultraviolet light, processed food, drugs (paracetamol, diclofenac, cisplatin, chlorpromazine, dexamethasone, etc.), various kinds of pollutants, etc., which lead to the generation of various free radicals (reactive molecules) such as ROS and Reactive Nitrogen Species (RNS). NOS, lipoxygenase, NADPH oxidase, mitochondrial Electron Transport Chain (ETC), and other naturally occurring enzymes aid in the production of different free radicals, including ROS and RNS. The mitochondrial ETC, NOS, NADPH oxidase, and lipoxygenase are examples of these endogenous sources (Figure 1).

Mitochondria are also responsible for the generation of many reactive molecules like ROS, besides essential cell organelles for ATP generation (Figure 2). Therefore, mitochondrial health is very important for maintaining spermatozoal motility and their forward progression during fertilization. The primary and major ROS is superoxide which is produced in mitochondria. Superoxide dismutase is an enzyme that helps to




Figure 1. Pathophysiology of oxidative stress-induced male and female infertility.




Figure 2. Pathophysiology of oxidative stress-induced mitochondrial dysfunction and spermatozoa damage in testes

convert harmful molecules of superoxide into hydrogen peroxide and molecular oxygen which ameliorates oxidative stress.

In NOS, Nitric Oxide (NO) is a principal factor released from normal endothelial and in inflammation, its production by the vasculature increases significantly and in combination with other ROS, contributes to oxidative stress. The plasma membrane contains the membrane-bound enzyme complex known as Nicotinamide Adenine

Dinucleotide Phosphate (NADPH) oxidase, which faces extracellular space. One of the main sources of cellular ROS is NADPH oxidases (NOXes), which continue to play a significant role in the production of ROS in normal conditions (Figures 2, 3). Lipid peroxidation is facilitated by lipoxygenases, which also catalyze the deoxygenation of polyunsaturated fatty acids in lipids and brain tissue^{25,26}. So, concurring with the present status of knowledge on oxidative stress and male reproduction, we suggest that

a heightened vulnerability to oxidative stress might be a common cause of infertility in males.

3. Oxidative Stress-Mediated Male Infertility in the High-Altitude Environment

Extreme climate, cold stress, high UV-radiation, and hypoxia in high altitude environments cumulatively affect the hypothalamus–hypophysis–gonad axis, and several other cellular and physiological mechanisms associated with maintaining oxygen homeostasis, cardiovascular, respiratory, and reproductive functions in human and animal²⁷⁻²⁹. Furthermore, hypoxia induces ROS generation, which later affects cellular components like protein, carbohydrates, lipids, and DNA³⁰. Since, spermatozoal integrity is maintained by cholesterol and lipid structure, which is significantly, affected under hypoxia-induced oxidative stress, spermatozoal damage is very common at high altitudes^{27,31,32}.

Several human and rodent studies reported imbalances in hormonal secretion, ovulatory activity, male infertility, and structural changes in testicular, ovarian, and prostate tissues due to hypoxia and oxygen deprivation at high altitudes²⁹⁻³⁷. Seminal examination revealed a significant reduction in sperm motility, volume, density, total counts, increased dimorphisms, and head damage in males exposed to hypoxia prevalent at high altitudes^{27,28,32-37}. Therefore, the likelihood of becoming pregnant is reduced in both people and animals that live at high elevations. It was reported that reduction in sperm concentration and testicular dysfunction can be the consequences of physical exercise at high altitudes³⁸. A recent study reported a decline in semen antioxidant potential in lowlander men presented to high altitude, whereas antioxidant activity was restored to some extent on 70 days onward arrival to the plain region³⁷. Another simulation study depicting hypoxia in male rats revealed significant damage to seminiferous tubules, reduction in testicular and epididymis weights, higher apoptosis in germ cells, and low sperm concentration²⁸. This damage to gonadal tissues also affects imbalance in reproductive hormones, viz. low levels of FSH, LH, testosterone, and estrogen. In conclusion, a high-altitude environment and hypoxia significantly affect the male reproductive functions. However, there are conflicting reports on the reversal of adverse changes in male reproductive

physiology and structural changes due to exposure to high altitude after coming to low altitude and plain areas. This needs to be investigated separately in the pre-pubertal, puberty, and post-pubertal stages considering the growth phase and reproductive stage of males during high-altitude exposure and post-induction. Therefore, environmental and physical stress at high altitudes limit the performance and reproduction in lowlander susceptible animals and human beings, which can be improved through antioxidant supplementation and exposure normoxic environment.

4. Mechanism of Oxidative Stress-Induced Spermatozoa Damage

Since spermatozoa are very dynamic and motile, they require for their activation a continuous supply of energy so that they are ironic in mitochondria¹⁹. Thus, if the semen has dysfunctional spermatozoa, it significantly raises the production of ROS, and whenever the ROS level surges it affects mitochondrial activity and, later, functions performed by sperm, such as motility. Recent studies have shown that there is leakage of an electron from actively respiring spermatozoa which generates mostly ROS which are free radicals³⁸. The ROS, such as hydroxyl (OH), superoxide (O₂⁻), nitric oxide (NO), peroxy (RO₂), lipid peroxy (LOO) and Thiol [RS-], and non-radical compounds (ozone [O₃], hydrogen peroxide [H₂O₂], lipid peroxide [LOOH], hypo chloric acid [HOCL], and singlet oxygen [-1O₂]), are generated which contain an unpaired electron that makes highly reactant to any biomolecules^{18,19,39}. Excessive semen ROS has been found anywhere from 30 to 80 percent of male infertile²⁴. Systems that produce ROS have also evidently demonstrated the vulnerability-bound hydrogen peroxide to be the most deadly oxygen metabolite concerning oxidative stress on sperm motility⁴⁰.

Within human spermatozoa, the majority of ROS is O₂⁻ (oxygen radical), and when O₂⁻ reacts with H₂ it leads to the production of H₂O₂, and thereby OH radicals are generated. Due to the disturbance of membrane fluidity, these OH-radicals are the strong initiators of the LPO cascade and can cause sperm function to be lost⁴¹⁻⁴³. Hence, sperm cells are prone to lipid peroxidation due to the high levels of unsaturated fatty acids they contain (Figure 3). Since the energy needed to dissociate carbon

and hydrogen is lowest at the mostly methylene position, the latter is more vulnerable to attack by free radicals. As of significance, the hydrogen constructs the process that starts lipid peroxidation get encouraged, producing a lipid radical with a carbon core that, when combined with oxygen, produces peroxy (ROO•) and alkoxy (RO•) radicals, which stabilize and eliminate hydrogen atoms from neighbouring carbons. Now, it is generally agreed upon that semen OS generates sperm cell membrane LPO, and male Sperm DNA Fragmentation (SDF), resulting in apoptosis²⁴. So, it is clear that in situations where motility is normal, oxidative stress can impair spermatozoa's ability to fertilize⁴⁴.

There are two major methods of free radical production in spermatozoa: - (i) Nicotinamide adenine dinucleotide phosphate oxidase system present in the sperm plasma membrane may lead to the production of ROS. (ii) In the sperm mitochondria, the nicotinamide adenine dinucleotide-dependent oxidoreductase reaction may also lead to the generation of reactive molecules (Figures 2, 3).

5. Gene Responsible for Abnormality of Spermatozoa and Male Infertility

The gene *NOX5* in the sperm encodes an enzyme NOX5, a calcium-dependent NADPH oxidase found in

the acrosomal and mid-piece regions which facilitates the production of O₂ in spermatozoa⁴⁵. NOX5 was first discovered in the human testis. It is activated by calcium binding to its N-terminal domain, which causes changes in cell shape and leads to oxidative stress. This finding provides further evidence that NOX5 is a major source of ROS generation in human spermatozoa. It is unknown if people with infertility linked to Oxidative Stress (OS) have increased NOX5 in their spermatozoa⁴¹. The morphologically aberrant, immature spermatozoa generate intracellular nicotinamide adenine dinucleotide phosphate (NADPH) and maintain an excess of the remnant body that contains significant levels of the cytoplasmic glucose-6-phosphate dehydrogenase enzyme. Afterwards, NADPH oxidase NOX5, which is intramembrane-based, converts NADPH to ROS.

Epigenetic modifications include DNA methylation, acetylation, phosphorylation, etc. It has been noted that oxidative stress can affect gene expression despite altering the order of genes, which has ramifications for epigenetics. ROS directly modify DNA, which influences male fertility. It has been demonstrated that disruption of DNA methylation disrupts spermatogenesis, resulting in a drop in the number of sperm as well as an upsurge in infertility^{46,47}. In addition, histone methylation dysregulation is linked to breaks in double-stranded




Figure 3. Oxidative stress-mediated spermatozoal dysfunction and male infertility

genetic material, with implications for infertility and the development of the next generation⁴⁸. Lastly, it is believed that microRNAs that control spermatogenesis in an epigenetic manner are critical for preserving fertility. A study was conducted on male patients with infertility in non-obstructive azoospermia, the levels of three specific microRNAs (miR713p, miR429, and miR141) were significantly increased⁴⁹. The epigenetic process linking oxidative stress to infertility is still under investigation. Once revealed, they provide a better understanding of molecular pathology, serve as therapeutic approaches, and afford insights regarding infertility components that are passed down from generation to generation⁵⁰.

6. Comparative Aspects of Oxidative Stress in Males and Females

The different results suggest that males might be more vulnerable to an increase in breeding effort, which is due to a decrease in antioxidant defence in comparison to females. ROS are essential for reproductive processes including ovulation, but when levels are too high, they can damage oocytes through oxidative stress and result in infertility. Melatonin's antioxidant action inside follicles may shield oocytes from ROS⁵¹. Reproductive processes including the growth of follicles, oocyte maturation, ovulation, fertilization, implantation, and the development of embryos are all impacted by ROS. In conjunction with the stimulation of vascularization within follicles, vascular endothelial cells and macrophages produce a significant quantity of ROS during the ovulation process following the Luteinizing Hormone (LH) surge⁵¹. The oxidative stress that ROS produced in follicles during ovulation causes in the oocyte and granulosa cells deteriorations of the integrity of the oocytes and may potentially lead to infertility⁵¹. The administration of melatonin may develop into a potential therapeutic approach to stop the aging-related decline in oocyte quantity and quality.

7. The Role of Food-Grade Antioxidants and Melatonin in Male Infertility

From various studies, it has been found that in a healthy male, two main mechanisms protect sperm DNA from OS. In the beginning, the genetic material is densely

coiled and packed with the histone proteins known as chromatin to facilitate that the genetic materials are slightly bare to free radicals⁵². Secondly, there are many natural antioxidants present in spermatozoa and seminal plasma which leads to minimizing ROS production and maintenance to its normal levels³⁸. These antioxidants prevent free radical formation by reacting with and then neutralizing them, which prevents OS onset and preserves the spermatozoa⁵³. Antioxidants including coenzyme Q10 and lactoferrin are also present in sperm cells⁵⁴. However, in a few infertile men, an insufficient number of antioxidants interferes with OS's complex stability and ends up in OS³⁸.

The body's antioxidant defence balance is crucial to keep the equilibrium among free radicals in addition to antioxidants; when it fails, the exogenous antioxidant supplements can play a preventive role. The foundation of oral antioxidant treatment is the understanding that initial oxidative damage develops because of increased ROS production and reduced seminal antioxidant status. Diet plays a very crucial role in day-to-day health benefits. So, diets rich in antioxidant contents can prevent the formation of oxidative stress in the testes so it must be consumed⁵⁴. The exogenous antioxidant category includes a large number of oral antioxidants, which the body receives from food sources, e.g., Vitamin C, Vitamin E, Coenzyme Q10, N acetylcysteine, carnitines, zinc, selenium, and pentoxifylline. Numerous research reports have been carried out to evaluate the efficacy of oral intake of antioxidants as a male infertility treatment.

Numerous physiological functions that are essential to our bodies, such as breathing, digestion, metabolizing alcohol, drug exposure, heat, environmental damage, pollutants, and metallic substances play a role in the development of OS, so all these have been shown to contribute to OS. It is reported that oxidative stress is associated with the development of numerous metabolic, and chronic disorders or cancers^{17,18,55}. It is also recognized, in addition, that activities that may cause a rise in the internal body temperature of the scrotal area such as lengthy, prolonged work hours, hot baths, gyms, and prolonged driving should be avoided. Lifestyle modifications can help to reduce the generation of ROS and perhaps address the disparities that lead to a state of oxidative stress.

7.1 Vitamin C and Carotenoids

Vitamin C and carotenoids are natural and essential antioxidants. They preserve the integrity of cell membranes

and control the process of spermatogenesis. They also play a role in regulating spermatogenesis. Citrus fruits and effervescent berries are good sources of water-soluble vitamin C, which has antioxidant qualities. Several studies have examined the effects of vitamin C consumption on spermatozoa and found noteworthy antioxidant benefits⁵⁶. Furthermore, decreased Vitamin C and expanded ROS ranges had been found in the reproductive fluid (seminal fluid) of males having azoospermia⁵⁷. Vegetable hues such as orange, red, yellow, and purple naturally include substances called carotenoids. These substances function as predecessors of vitamin A, of which retinol is a necessary element. Carotenoid deficiency can result in reduced sperm motility and male infertility⁵⁸.

7.2 Vitamin E

Vitamin E (alpha-tocopherol) is a biological compound that is fat-soluble and frequently distributed across cell membranes (Table 1). Its antioxidant activity is based primarily on suppressing ROS-induced peroxidation of lipids and removing the hydroxyl extra electron-containing molecule (free radicals) and superoxide. Therefore, alpha-tocopherol primarily offers protection from free radical damage in the components of the sperm membrane and, to a lesser extent, reduces ROS production. Therefore, Vitamin E primarily protects sperm membrane components from damage and, to a lesser extent, reduces ROS production. Its oral administration significantly increases sperm motility by reducing the malondialdehyde an end product of lipid peroxidation at the time of sperm

production and is, therefore, a secondary marker of the intensity of intracellular processes⁵⁹.

7.3 Carnitine

Carnitine or levocarnitine is a naturally occurring amino acid that plays a role in mitochondrial transfer and oxidation of long-chain fatty acids, its chemical structure was established in 1927 as (3-hydroxy-4- trimethylamine butyrate)^{60,61}. It serves as a vital substance that helps with the transportation of long-chain fatty acids within the mitochondrial matrix. This process facilitates cellular energy production through oxidation^{62,63}. An important aspect here is the high levels of carnitine in the male reproductive tract, particularly the epididymis. This suggests that it plays an important function in the consumption of energy and maturation of sperm^{64,65}.

7.4 Melatonin

Oxidative stress-induced endocrine disruption has a deleterious effect on pineal gland secretion which affects the reduction in melatonin levels in the body⁶⁶. Thereafter, it has a chain of effects on the biological clock and metabolic function. Melatonin, a neurohormone, is an excellent free-radical scavenger. It scavenges/neutralizes free radicals (ROS) which are produced by oxidative stress. Melatonin also activates the major enzymatic antioxidant mechanisms of body tissues^{67,68} and is also found in human sperm⁶⁹. Melatonin, with its antioxidant and anti-apoptotic effects, up-regulates antioxidant defence mechanisms and inhibits apoptosis, thus preventing dexamethasone-

Table 1. Physiological and ameliorative role of some food-grade natural antioxidants in male reproduction and fertility

Sl. No.	Name of Antioxidant	Role and their action	Effects
1	Vitamin C	Counterbalance free radicals	Protects viability as well as motility.
2	Vitamin E	Neutralizes free radicals	Enhances the function /activity of many antioxidants and reduces the oxidation of lipids.
3	Cysteines	Reduced glutathione (GSH) formation is increased by cysteines	Prevent lipid peroxidation.
4	Pentoxifylline	An important second messenger cAMP breakdown is prevented and it conquers the formation of pro-inflammatory factors	Reduces the peroxidation of lipids
5	Carotenoids	Quenche singlet molecular oxygen	Limit the oxidation of lipids
6	Carnitine	An energy source and balance between free radicals and antioxidants by reducing reactive molecules (free radicals)	Prevents lipid peroxidation and also protects from damage to the DNA
7	Bilirubin	Extracellular antioxidants	

induced testicular damage⁷⁰. In addition to eliminating the very harmful hydroxyl radical (OH)^{71,72}, it has been reported that melatonin also eliminates many reactive oxygen and nitrogen species such as singlet oxygen ($1O_2$)⁷³, nitric oxide (NO)⁷⁴, peroxyxynitrite anion (ONOO) and/or its metabolites⁷⁴, and hydrogen peroxide (H_2O_2)⁷⁵. In addition to these, melatonin changes the enzyme activities which metabolize free radicals⁷⁶⁻⁷⁸ and alters membrane fluidity^{79,80}. This reduces the ability of damaging species to attack macromolecules in this structure.

8. Conclusion

Oxidative stress and free radicals are known to be detrimental to the health of a healthy individual. This review indicates that oxidative stress is central to structural and DNA damage of spermatozoa and induces male infertility in plain and high-altitude regions. Therefore, the evaluation of antioxidant enzymes, oxidative stress parameters, and body antioxidant levels are necessary for new diagnostic methods for diagnosing diseases related to reproduction. In addition to assessing sperm oxidative stress, blood redox status, and leukocyte ROS level, this also opens new possibilities and small applications for specialists to test sperm quality and monitor fertility. There are various antioxidant supplements available, including melatonin, which may be used in male infertility to ameliorate free radicals-mediated/induced male infertility. Therefore, dealing with the redox parameters may help reduce systemic oxidative stress as an aspect of male infertility. It can help develop novel treatment strategies by taking antioxidant supplements to improve the detection along management of male infertility.

9. Acknowledgment

All authors thank the Department of Zoology, Guru Ghasidas Vishwavidyalaya, Bilaspur, Chhattisgarh, and the Department of Advanced Research (DIHAR), DRDO, Leh, India, for providing support and space. The first author (SP) is also grateful to the University Grants Commission (Selection and Award Bureau), New Delhi, for providing the Rajiv Gandhi National Fellowship for Other Backward Classes (RGNFOBC) during this study.

10. References

1. Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *Reprod Biol* Endocrinol. 2015; 13(1):1-9. <https://doi.org/10.1186/s12958-015-0032-1> PMID: 25928197 PMCID: PMC4424520
2. Baskaran S, Finelli R, Agarwal A, Henkel R. Diagnostic value of routine semen analysis in clinical andrology. *Andrologia*. 2021; 53(2). <https://doi.org/10.1111/and.13614> PMID: 32400107
3. Slade P, O'Neill C, Simpson AJ, Lashen H. The relationship between perceived stigma, disclosure patterns, support, and distress in new attendees at an infertility clinic. *Hum Reprod*. 2007; 22:2309-7. <https://doi.org/10.1093/humrep/dem115> PMID: 17580298
4. Stearns SC. *The evolution of life histories*, Vol. 249. Oxford: Oxford University Press; 1992. p.xii
5. Zera AJ, Harshman LG. The physiology of life history trade-offs in animals. *Ann Rev Ecol Syst*. 2001; 32(1):95-126. <https://doi.org/10.1146/annurev.ecolsys.32.081501.114006>
6. Alvarez AC, Bertrand S, Devevey G, Prost J, Faivre B, Sorci G. Increased susceptibility to oxidative stress as a proximate cost of reproduction. *Ecol Lett*. 2004; 7(5):363-8. <https://doi.org/10.1111/j.1461-0248.2004.00594.x>
7. Johnston, SL, Souter DM, Erwin SS, Tolkamp BJ, Yearsley JM, Gordon IJ, *et al*. Associations between basal metabolic rate and reproductive performance in C57BL/6J mice. *J Exp Biol*. 2007; 210(1):65-74 <https://doi.org/10.1242/jeb.02625> PMID: 17170149.
8. Hedger MP. The immunophysiology of male reproduction. *Knobil and Neill's Physiology of Reproduction*. 2015; 805-92. <https://doi.org/10.1016/B978-0-12-397175-3.00019-3>
9. Ye L, Huang W, Liu S, Cai S, Hong L, Xiao W, *et al*. Impacts of immunometabolism on male reproduction. *Front Immunol*. 2021; 12. <https://doi.org/10.3389/fimmu.2021.658432> PMID: 34367130, PMCID: PMC8334851
10. Angilletta MJ, Sears MW. The metabolic cost of reproduction in an oviparous lizard. *Funct Ecol*. 2000; 14(1):39-45. <https://doi.org/10.1046/j.1365-2435.2000.00387.x>
11. Kohen R, Nyska A. Invited review: Oxidation of biological systems: Oxidative stress phenomena, antioxidants, redox reactions, and methods for their quantification. *Toxicol Pathol*. 2002; 30(6):620-50. <https://doi.org/10.1080/01926230290166724> PMID: 12512863
12. Aitken RJ, Curry BJ. Redox regulation of human sperm function: From the physiological control of sperm capacitation to the etiology of infertility and DNA damage in the germ line. *Antioxid Redox Signal*. 2011; 14:367-81. <https://doi.org/10.1089/ars.2010.3186> PMID: 20522002
13. Aitken RJ, Gibb Z, Mitchell LA, Lambourne SR, Connaughton HS, De Iulius GN. Sperm motility is lost in vitro as a consequence of mitochondrial free radical production and the generation of electrophilic aldehydes but can be significantly rescued by the presence of nucleophilic thiols. *Biol Reprod*. 2012; 87. <https://doi.org/10.1095/biolreprod.112.102020> PMID: 22933515

14. Garratt M, Brooks RC. Oxidative stress and condition-dependent sexual signals: More than just seeing red. *Proc Royal Soc B: Biol Sci.* 2012; 279(1741):3121-30. <https://doi.org/10.1098/rspb.2012.0568> PMID: 22648155, PMCID: PMC3385731
15. Sies H, Jones DP. Reactive Oxygen Species (ROS) as pleiotropic physiological signalling agents. *Nat Rev Mol Cell Biol.* 2020; 21(7):363-83. <https://doi.org/10.1038/s41580-020-0230-3> PMID: 32231263
16. Tan BL, Norhaizan ME, Liew WPP. Nutrients and oxidative stress: Friend or foe? *Oxid Med Cell Longev.* 2018. <https://doi.org/10.1155/2018/9719584> PMID: 29643982 PMCID: PMC5831951
17. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. *Nature.* 2000; 408(6809):239- 47. <https://doi.org/10.1038/35041687> PMID: 11089981
18. Miranda-Vilela AL, Alves PC, Akimoto AK, Pereira LCS, Klautau-Guimarães Md, Grisolia CK. The effect of hydrogen peroxide-induced oxidative stress on leukocytes depends on age and physical training in healthy human subjects carrying the same genotypes of antioxidant enzymes' gene polymorphisms. *Am J Hum Biol.* 2010; 22:807-12. <https://doi.org/10.1002/ajhb.21086> PMID: 20878976
19. Henkel RR. Leukocytes and oxidative stress: Dilemma for sperm function and male fertility. *Asian J Androl.* 2011; 13:43-52. <https://doi.org/10.1038/aja.2010.76> PMID: 21076433 PMCID: PMC3739401
20. Surai PF. Selenium in poultry nutrition 1. Antioxidant properties, deficiency and toxicity. *Poult Sci J.* 2002; 58(3):333-47. <https://doi.org/10.1079/WPS20020026>
21. Sharma RK, Pasqualotto FF, Nelson Thomas AJ, Agarwal A. Relationship between seminal white blood cell counts and oxidative stress in men treated at an infertility clinic. *J Androl.* 2001; 22(4):575-83. <https://doi.org/10.1002/j.1939-4640.2001.tb02217.x> PMID: 11451354
22. Koppers AJ, Garg ML, Aitken RJ. Stimulation of mitochondrial reactive oxygen species production by unesterified, unsaturated fatty acids in defective human spermatozoa. *Free Radic Biol Med.* 2010; 48(1):112-9. <https://doi.org/10.1016/j.freeradbiomed.2009.10.033> PMID: 19837155
23. Iwasaki A, Gagnon C. Formation of reactive oxygen species in spermatozoa of infertile patients. *Fertil Steril.* 1992; 57(2):409-16. [https://doi.org/10.1016/S0015-0282\(16\)54855-9](https://doi.org/10.1016/S0015-0282(16)54855-9) PMID: 1735495
24. Takeshima T, Kuroda S, Yumura Y. Alternations of sperm protein profiles to elucidate the mechanism of impaired spermatogenesis by cancer chemotherapy. *Fertil Steril.* 2020; 114(3):e256-7. <https://doi.org/10.1016/j.fertnstert.2020.10.022>
25. Czubowicz K, Czapski GA, Cieřlik M, Strosznajder RP. Lipoxigenase inhibitors protect brain cortex macromolecules against oxidation evoked by nitrosative stress. *Folia Neuropathol.* 2010; 48:289-92. PMID: 21225511
26. Czapski GA, Czubowicz K, Strosznajder RP. Evaluation of the antioxidative properties of lipoxigenase inhibitors. *Pharmacol Rep.* 2012; 64(5):1179-88. [https://doi.org/10.1016/S1734-1140\(12\)70914-3](https://doi.org/10.1016/S1734-1140(12)70914-3) PMID: 2323
27. Cofré E, Peralta OA, Raggi A, los Reyes MD, Sales F, González-Bulnes A, *et al.* Ram semen deterioration by short-term exposure to high altitude is prevented by improvement of antioxidant status. *Animal.* 2018; 12(5):1007-14. <https://doi.org/10.1017/S1751731117002452> PMID: 28988559
28. Li X-Y, Zhang M-H, Chen Z-W, Chen Z-W. Male reproductive system and simulated high-altitude environment: Preliminary results in rats. *Asian J Androl.* 2023; 25(3):426-32. PMID: 36510858 <https://doi.org/10.4103/aja202290> PMCID: PMC10226506
29. Kumari P, Bharti VK, Kumar K, Sharma I. Hypobaric hypoxia affects the reproductive physiology of dairy cattle. *Anim Reprod Update.* 2023; 3(2):6-17. <https://doi.org/10.48165/aru.2023.3.2.2>
30. Zepeda AB, Figueroa CA, Calaf GM, Farias JG. Male reproductive system and antioxidants in oxidative stress induced by hypobaric hypoxia. *Andrologia.* 2014; 46:1-8. <https://doi.org/10.1111/and.12039> PMID: 23127143
31. Cassinelli LM. Biology of fertility; acclimatization of the ovine animals to various altitudes. *An Univ nac Mayor de San Marcos.* 1953; 36:419-46. PMID: 13197786
32. Farias JG, Bustos-Obregón E, Orellana R, Bucarey JL, Quiroz E, Reyes JG. Effects of chronic hypobaric hypoxia on testis histology and round spermatid oxidative metabolism. *Andrologia.* 2005; 37:47-52 <https://doi.org/10.1111/j.1439-0272.2004.00651.x> PMID: 15644063
33. Saxena DK. Effect of hypoxia by intermittent altitude exposure on semen characteristics and testicular morphology of male rhesus monkeys. *Int J Biometeorol.* 1995; 38(3):137-40. <https://doi.org/10.1007/BF01208490> PMID: 7744528
34. Okumura A, Fuse H, Kawauchi Y, Mizuno I, Akashi T. Changes in male reproductive function after high altitude mountaineering. *High Alt Med Biol.* 2004; 4(3):349-53. <https://doi.org/10.1089/152702903769192304> PMID: 14561240
35. He J, Cui J, Wang R, Gao L, Gao X, Yang L, *et al.* Exposure to hypoxia at high altitude (5380 m) for 1 year induces reversible effects on semen quality and serum reproductive hormone levels in young male adults. *High Alt Med Biol.* 2015; 16(3):216-22. <https://doi.org/10.1089/ham.2014.1046> PMID: 26288097
36. Zheng S, Liu Y, Li P, Tian H. Short-term high-altitude exposure (3600 m) alters the type distribution of sperm deformity. *High Alt Med Biol.* 2019; 20(2):198-202. <https://doi.org/10.1089/ham.2018.0133> PMID: 30900912

37. Verratti V, Mrakic-Spota S, Fusi J, Sabovic I, Franzoni F, Pietrangelo T, *et al.* Fertility impairment after trekking at high altitude: A proof of mechanisms on redox and metabolic seminal changes. *Int J Mol Sci.* 2022; 23(16). <https://doi.org/10.3390/ijms23169066> PMID: 36012330, PMCID: PMC9409093
38. Agarwal A, Virk G, Ong C, Du Plessiss SS. Effect of oxidative stress on male reproduction. *World J Mens Health.* 2014; 32(1):1-17. <https://doi.org/10.5534/wjmh.2014.32.1.1> PMID: 24872947, PMCID: PMC4026229
39. Bansal AK, Bilaspuri GS. Impacts of oxidative stress and antioxidants on semen functions. *Vet Med Int.* 2010. <https://doi.org/10.4061/2011/686137> PMID: 20871827, PMCID: PMC2943128
40. Baumber J, Ball BA, Gravance CG, Medina V, Davies-Morel MCG. The effect of reactive oxygen species on equine sperm motility, viability, acrosomal integrity, mitochondrial membrane potential, and membrane lipid peroxidation. *J Androl.* 2000; 21(6): 895-902. <https://doi.org/10.1002/j.1939-4640.2000.tb03420.x>
41. Chen SJ, Allam JP, Duan YG, Haidl G. Influence of reactive oxygen species on human sperm functions and fertilizing capacity including therapeutical approaches. *Arch Gynecol Obstet.* 2013; 288(1):191-9. <https://doi.org/10.1007/s00404-013-2801-4> PMID: 23543240
42. Hazout A, Menezo Y, Madelenat P, Yazbeck C, Selva J, Cohen-Bacrie P. Causes and clinical implications of sperm DNA damages. *Gynecol Obstet Fertil.* 2008; 36(11):1109-17. <https://doi.org/10.1016/j.gyobfe.2008.07.017> PMID: 18964175
43. Sikka SC. Relative impact of oxidative stress on male reproductive function. *Curr Med Chem.* 2001; 8(7):851-62. <https://doi.org/10.2174/0929867013373039> PMID: 11375755
44. Wishart GJ. Effects of lipid peroxide formation in fowl semen on sperm motility, ATP content and fertilizing ability. *Reproduction.* 1984; 71(1):113-8. <https://doi.org/10.1530/jrf.0.0710113> PMID: 6726671
45. Sabeur K, Ball BA. Characterization of NADPH oxidase 5 in equine testis and spermatozoa. *Reproduction.* 2007; 134(2):263-70. <https://doi.org/10.1530/REP-06-0120> PMID: 17660236
46. Laqqan M, Solomayer EF, Hammadeh M. Aberrations in sperm DNA methylation patterns are associated with abnormalities in semen parameters of subfertile males. *Reprod Biol.* 2017; 17(3):246-51. <https://doi.org/10.1016/j.repbio.2017.05.010> PMID: 28566207
47. Urdinguio RG, Bayón GF, Dmitrijeva M, Torano EG. Aberrant DNA methylation patterns of spermatozoa in men with unexplained infertility. *Hum Reprod.* 2015; 30(5):1014-28. <https://doi.org/10.1093/humrep/dev053> PMID: 25753583
48. Sharma U, Conine CC, Shea JM, Boskovic A. Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science.* 2016; 351(6271):391-96. <https://doi.org/10.1126/science.aad6780> PMID: 26721685 PMCID: PMC4888079
49. Hilz S, Modzelewski AJ, Cohen PE, Grimson A. The roles of microRNAs and siRNAs in mammalian spermatogenesis. *Development.* 2016; 143(17):3061-73. <https://doi.org/10.1242/dev.136721> PMID: 27578177 PMCID: PMC5047671
50. Ritchie C, Edmund YK. Oxidative stress in the pathophysiology of male infertility. *Andrologia.* 2021; 53(1). <https://doi.org/10.1111/and.13581> PMID: 32323352
51. Tamura H, Tanabe M, Jozaki M, Taketani T. Antioxidative action of melatonin and reproduction. *Glycative Stress Research.* 2019; 6(3):192-7.
52. Lampiao F. Free radicals generation in an *in vitro* fertilization setting and how to minimize them. *World J Obstet Gynecol.* 2012; 1(3):29-34. <https://doi.org/10.5317/wjog.v1.i3.29>
53. Sharma, RK, Agarwal A. Role of reactive oxygen species in male infertility. *Urology.* 1996; 48(6):835-50. [https://doi.org/10.1016/S0090-4295\(96\)00313-5](https://doi.org/10.1016/S0090-4295(96)00313-5) PMID: 8973665
54. Lanzafame FM, La Vignera S, Vicari E, Calogero AE. Oxidative stress and medical antioxidant treatment in male infertility. *Reprod Biomed Online.* 2009; 19(5):638-59. <https://doi.org/10.1016/j.rbmo.2009.09.014> PMID:20021713
55. Aminjan HH, Abtahi SR, Hazrati E, Chamanara M. Targeting of oxidative stress and inflammation through ROS/NF-kappaB pathway in phosphine-induced hepatotoxicity mitigation. *Life Sci.* 2019; 232. <https://doi.org/10.1016/j.lfs.2019.116607> PMID: 31254582
56. Das UB, Mallick M, Debnath JM, Ghosh D. Protective effect of ascorbic acid on cyclophosphamide – Induced testicular gametogenic and androgenic disorders in male rats. *Asian J Androl.* 2002; 4(3):201-7. PMID: 12364977
57. Lewis SE, Simon L. Clinical implications of sperm DNA damage. *Hum Fertil (Camb).* 2010; 13(4):201-7. <https://doi.org/10.3109/14647273.2010.528823> PMID: 21117929
58. Hogarth, CA, Griswold MD. The key role of Vitamin A in spermatogenesis. *J Clin Invest.* 2010; 120(4):956- 62. <https://doi.org/10.1172/JCI41303> PMID: 20364093 PMCID: PMC2846058
59. Suleiman SA, Ali ME, Zaki ZM. Lipid peroxidation and human sperm motility: Protective role of vitamin E. *J Androl.* 1996; 17(5):530-7. <https://doi.org/10.1002/j.1939-4640.1996.tb01830.x> PMID: 8957697
60. Kelly GS. L-Carnitine: Therapeutic applications of a conditionally-essential amino acid. *Altern Med Rev.* 1998; 3(5):345-60. PMID:9804680
61. Modanloo M, Shokrzadeh M. Analyzing mitochondrial dysfunction, oxidative stress, and apoptosis: Potential

- role of L-carnitine. *Iran J Kidney Dis.* 2019; 13(2). PMID: 30988244
62. Agarwal A, Said TM. Carnitines and male infertility. *Reprod Biomed Online.* 2004; 8(4):376-84. [https://doi.org/10.1016/S1472-6483\(10\)60920-0](https://doi.org/10.1016/S1472-6483(10)60920-0) PMID: 15149558
 63. Ng CM, Blackman MR, Wang C, Swerdloff RS. The role of carnitine in the male reproductive system. *Ann NY Acad Sci.* 2004; 1033(1):177-88. <https://doi.org/10.1196/annals.1320.017> PMID: 15591015
 64. Vicari E, Rubino C, De Palma, Longo G. Antioxidant therapeutic efficiency after the use of carnitine in infertile patients with bacterial or non-bacterialprostatitis-epididymitis. *Arch Ital Urol Androl.* 2001; 73(1):15-25. PMID: 11505809
 65. Mongioi L, Calogero AE, Vicari E, Condorelli RA. The role of carnitine in male infertility. *Andrology.* 2016; 4(5):800-7. <https://doi.org/10.1111/andr.12191> PMID: 27152678
 66. Aulinas A. Physiology of the pineal gland and melatonin. *Endotext [Internet].* 2019. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK550972/>
 67. Mayo JC, Sainz RM, Antoli I, Herrera F. Melatonin regulation of antioxidant enzyme gene expression. *Cell Mol Life Sci.* 2002; 59(10):1706-13. <https://doi.org/10.1007/PL00012498> PMID: 12475181
 68. Bharti VK, Srivastava RS. Pineal proteins up-regulate specific antioxidant defense systems in the brain. *Oxid Med Cell Longev.* 2009; 2(2):88-92. <https://doi.org/10.4161/oxim.2.2.8361>
 69. Awad H, Halawa F, Mostafa T, Atta H. Melatonin hormone profile in infertile males. *Int J Androl.* 2006; 29(3):409-13. <https://doi.org/10.1111/j.1365-2605.2005.00624.x> PMID: 16371109
 70. Mukherjee S, David A, Yadav S, Baluska F. Salt stress-induced seedling growth inhibition coincides with differential distribution of serotonin and melatonin in sunflower seedling roots and cotyledons. *Physiol Plant.* 2014; 152(4):714-28. <https://doi.org/10.1111/ppl.12218> PMID: 24799301
 71. Stasica P, Ulanski P, Rosiak JM. Melatonin as a hydroxyl radical scavenger. *J Pineal Res.* 1998; 25(1):65-6. <https://doi.org/10.1111/j.1600-079X.1998.tb00387.x> PMID: 9694406
 72. Tan DX, Manchester LC, Reiter RJ, Qi WB. Significance of melatonin in antioxidative defense system: Reactions and products. *Biol Signals Recept.* 2000; 9(3-4):137-59. <https://doi.org/10.1159/000014635> PMID: 10899700
 73. Zang LY, Cosma G, Gardner H, Vallyathan V. Scavenging of reactive oxygen species by melatonin. *Biochim Biophys Acta-Gen Sub.* 1998; 1425(3):469-77. [https://doi.org/10.1016/S0304-4165\(98\)00099-3](https://doi.org/10.1016/S0304-4165(98)00099-3) PMID: 9838210
 74. Noda Y, Mori A, Liburdy R, Packer L. Melatonin and its precursors scavenge nitric oxide. *J Pineal Res.* 1999; 27(3):159-63. <https://doi.org/10.1111/j.1600-079X.1999.tb00611.x> PMID: 10535765
 75. Zhang H, Squadrito GL, Uppu R, Pryor WA. Reaction of peroxyxynitrite with melatonin: A mechanistic study. *Chem Res Toxicol.* 1999; 12(6):526-34. <https://doi.org/10.1021/tx980243t> PMID: 10368316
 76. Pablos MI, Guerrero JM, Ortiz GG, Agapito M. Both melatonin and a putative nuclear receptor agonist CGP 52608 stimulate glutathione peroxidase and glutathione reductase activities in mouse brain in vivo. *Neuroendocrinol Lett.* 1997; 18:49-58.
 77. Reiter RJ, Tang L, Garcia JJ, Muñoz-Hoyos A. Pharmacological actions of melatonin in oxygen radical pathophysiology. *Life Sci.* 1997; 60(25):2255-71. [https://doi.org/10.1016/S0024-3205\(97\)00030-1](https://doi.org/10.1016/S0024-3205(97)00030-1) PMID: 9194681
 78. Hara M, Iigo M, Ohtani-Kaneko R, Nakamura N. Administration of melatonin and related indoles prevents exercise-induced cellular oxidative changes in rats. *Biol Signals.* 1997; 6(2):90-100. <https://doi.org/10.1159/000109113> PMID: 9266096
 79. Garcia JJ, Reiter RJ, Guerrero JM, Escames G. Melatonin prevents changes in microsomal membrane fluidity during induced lipid peroxidation. *FEBS Lett.* 1997; 408(3):297-300. [https://doi.org/10.1016/S0014-5793\(97\)00447-X](https://doi.org/10.1016/S0014-5793(97)00447-X) PMID: 9188780
 80. Garcia JJ, Reiter RJ, Ortiz GG, Oh CS. Melatonin enhances tamoxifen's ability to prevent the reduction in microsomal membrane fluidity induced by lipid peroxidation. *J Membr Biol.* 1998; 162(1):59-65. <https://doi.org/10.1007/s002329900342> PMID: 9516238